

Supplementary Table 2. CLIF-SOFA variables as risk factors for 28-day mortality in all patients with variceal bleeding (n=474)*

Variable	Univariate analysis		Multivariate analysis	
	HR (95% CI)	P-value	HR (95% CI)	P-value
Bilirubin (mg/dL)	1.10 (1.07–1.13)	<0.001	1.03 (0.99–1.07)	0.152
Creatinine (mg/dL)	1.43 (1.27–1.60)	<0.001	1.29 (1.09–1.53)	0.003
HE (presence)	6.57 (3.02–14.31)	<0.001	3.30 (1.39–7.87)	0.007
Prothrombin time (INR)	3.19 (2.48–4.10)	<0.001	2.20 (1.62–3.00)	<0.001
MAP (<70 mmHg)	1.50 (0.63–3.59)	0.358		
PaO ₂ /FiO ₂ (≤400)	1.20 (0.67–3.07)	0.705		

CLIF SOFA, the chronic liver failure-sequential organ failure assessment; HR, hazard ratio; CI, confidence interval; HE, hepatic encephalopathy; INR, international ratio; MAP, mean arterial pressure; PaO₂, partial pressure of arterial oxygen; FiO₂, fraction of inspired oxygen.

*Subjects (n=474), event: death (n=39).