

Supplementary Figure 7. Protein quantitation for western blotting. (A) Quantification of GOLM1, OPN, and ABCG5 protein levels in HepG2-shGOLM1-1/-2 cells. (B) Quantification of GOLM1, OPN, and ABCG5 protein levels in HepG2-GOLM1 cells. (C) Quantification of GOLM1, OPN, and ABCG5 protein levels in HepG2-shOPN-1/-2 cells. (D) Quantification of GOLM1, OPN, and ABCG5 protein levels in HepG2-OPN cells. (E) Quantification of GOLM1, OPN, and ABCG5 protein levels in shGOLM1, shGOLM1+OPN-OE, GOLM1-OE, and GOLM1-OE+shOPN HepG2 cells. ns, nonsignificant; ***P<0.005. ABCG5, ATP binding cassette transporter subfamily G member 5; GOLM1, Golgi membrane protein 1; HepG2-GOLM1 cells, GOLM1-overexpressing HepG2 cells; HepG2-OPN cells, OPN-overexpressing HepG2 cells; HepG2-shGOLM1 cells, GOLM1-knockdown HepG2 cells; HepG2-shOPN cells, OPN-knockdown HepG2 cells; OPN, osteopontin.

 https://doi.org/10.3350/cmh.2024.0657